


Anti-PD-1 | Pembrolizumab Biosimilar  
Anti-PD-L1 | Atezolizumab Biosimilar  
Anti-CTLA-4 | Ipilimumab Biosimilar  
Anti-HER2 | Trastuzumab Biosimilar  
Anti-EGFR | Cetuximab Biosimilar  
and more



## The Journal of Immunology

RESEARCH ARTICLE | MARCH 15 2012

### Chronic Shift-Lag Alters the Circadian Clock of NK Cells and Promotes Lung Cancer Growth in Rats **FREE**

Ryan W. Logan; ... et. al

*J Immunol* (2012) 188 (6): 2583–2591.

<https://doi.org/10.4049/jimmunol.1102715>

#### Related Content

Suprachiasmatic Astrocytes Modulate the Circadian Clock in Response to TNF- $\alpha$

*J Immunol* (November,2013)

Dysregulation of Inflammatory Responses by Chronic Circadian Disruption

*J Immunol* (November,2010)

Circadian Oscillations of Clock Genes, Cytolytic Factors, and Cytokines in Rat NK Cells

*J Immunol* (June,2005)

# Chronic Shift-Lag Alters the Circadian Clock of NK Cells and Promotes Lung Cancer Growth in Rats

Ryan W. Logan,\* Changqing Zhang,\* Sengottuvelan Murugan,\* Stephanie O'Connell,\* Dale Levitt,\* Alan M. Rosenwasser,<sup>†</sup> and Dipak K. Sarkar\*

Prolonged subjection to unstable work or lighting schedules, particularly in rotating shift-workers, is associated with an increased risk of immune-related diseases, including several cancers. Consequences of chronic circadian disruption may also extend to the innate immune system to promote cancer growth, as NK cell function is modulated by circadian mechanisms and plays a key role in lysis of tumor cells. To determine if NK cell function is disrupted by a model of human shift-work and jet-lag, Fischer (344) rats were exposed to either a standard 12:12 light-dark cycle or a chronic shift-lag paradigm consisting of 10 repeated 6-h photic advances occurring every 2 d, followed by 5–7 d of constant darkness. This model resulted in considerable circadian disruption, as assessed by circadian running-wheel activity. NK cells were enriched from control and shifted animals, and gene, protein, and cytolytic activity assays were performed. Chronic shift-lag altered the circadian expression of clock genes, *Per2* and *Bmal1*, and cytolytic factors, perforin and granzyme B, as well as the cytokine, *IFN- $\gamma$* . These alterations were correlated with suppressed circadian expression of NK cytolytic activity. Further, chronic shift-lag attenuated NK cell cytolytic activity under stimulated *in vivo* conditions, and promoted lung tumor growth following *i.v.* injection of MADB106 tumor cells. Together, these findings suggest chronic circadian disruption promotes tumor growth by altering the circadian rhythms of NK cell function. *The Journal of Immunology*, 2012, 188: 2583–2591.

The multioscillator circadian system adapts to changing internal and external states to optimize the timing of physiological processes. Temporal coordination among multiple physiological systems is essential for homeostatic regulation, whereas circadian disruption may negatively impact health. Several large-scale epidemiological studies on rotating shift-workers have reported that working during the night is a major risk factor for several types of cancer, including non-Hodgkin's lymphoma (1) and breast (2–4), endometrial (5), prostate (6, 7), and colon (8, 9) cancers. Further, cancer risk is positively related to the frequency of night shifts an individual had worked, highlighting the potential detrimental effects of prolonged unstable work and light-exposure schedules (4, 9). In addition, robustness of circadian rhythms in salivary cortisol levels and rest-activity cycles are significant predictors for survival in cancer patients (10, 11). Other diseases, such as obesity, diabetes, and cardiovascular problems, are highly prevalent among shift-workers as well (12–14). Despite growing evidence from human studies linking circadian disruption, or desynchrony, to disease, an understanding of the essential mechanisms involved in the transition or promotion of particular disease states, such as tumor development, is lacking.

Since the initial reports associating shift-work and cancer in humans, a majority of animal studies have primarily focused on the effects of disrupted molecular clocks on cellular proliferation pathways to promote tumor growth (15–19). Although these pathways are important for cancer development and progression, the relationship between circadian disruption and cancer may extend beyond perturbations of cell cycle processes. For example, the inflammatory response by macrophages is dysregulated by disruption to cellular clocks (20, 21). However, the underlying mechanism by which altered circadian immune function may promote the transition to particular disease states is unknown, especially in relation to cancer. There is little evidence implicating disruption to circadian mechanisms in impaired immune function, as well as tumor growth.

The role of NK cells as critical mediators of cancer immunosurveillance is well established (22, 23). In mice, depletion of NK cells is associated with increased tumor growth in spontaneous and induced tumor models (24, 25). Similarly, low NK cell activity is associated with increased risk for cancer in human populations (26, 27). To kill tumor cells, NK cells recruit pro-inflammatory cytokines, release cytolytic granules, and activate receptors on target cells (23, 28, 29). Of particular importance are the cytolytic factors granzyme B and perforin and cytokines TNF and *IFN- $\gamma$* , which are critical factors involved in regulating NK cell-mediated killing of tumor cells (28–30). Decreases in any of these factors are associated with increased risk of developing infections and tumors (24, 25, 31–33).

The negative health consequences of circadian disruption are likely to involve alterations in NK cell function. Circadian mechanisms may optimize an immunological response by coordinating function between peripheral tissues and immunocompetent cells. Previously, we have reported that NK cell function, including cytokines, cytolytic factors, and cytolytic ability, is tightly controlled by the circadian system (34). Signals directly or indirectly derived from the circadian pacemaker, which is located

\*Endocrine Program, Rutgers, The State University of New Jersey, New Brunswick, NJ 08901; and <sup>†</sup>Department of Psychology, University of Maine, Orono, ME 04469  
Received for publication October 13, 2011. Accepted for publication January 2, 2012.

This work was supported by National Institutes of Health Grant R01 HL088041 and R37 Award AA08757.

Address correspondence and reprint requests to Prof. Dipak K. Sarkar, Director, Endocrine Program, Department of Animal Sciences, Rutgers, The State University of New Jersey, 67 Poultry Farm Lane, New Brunswick, NJ 08901. E-mail address: sarkar@aesop.rutgers.edu

Abbreviations used in this article: CT, circadian time; DD, total darkness; LD, light-dark (cycle); SCN, suprachiasmatic nucleus.

Copyright © 2012 by The American Association of Immunologists, Inc. 0022-1767/12/\$16.00

in the suprachiasmatic nucleus (SCN) of the anterior hypothalamus, are transmitted to peripheral tissues by neural and endocrine pathways to synchronize and coordinate body physiology (34–36). Rhythmic sympathetic input to the spleen modulates NK cell function by regulating components of the molecular clock in these cells (37). Maximal cytotoxicity at specific circadian phases is postulated to be driven by molecular clock-coordinated expression of IFN- $\gamma$ , TNF- $\alpha$ , perforin, and granzyme B (38, 39).

Nevertheless, few data indicating potential relationships between the disruption of clocks in immune cells and the promotion of tumor growth are available. Therefore, the main goal of this study was to determine whether NK cells were susceptible to circadian disruption by exposing animals to a chronic shift-lag paradigm in which the daily light-dark (LD) cycle was repeatedly phase shifted to experimentally mimic the disruptive effects of human shift-work and jet-lag. Under the shift-lag schedule, circadian disruption was assessed by continuously monitoring voluntary running-wheel activity. We then examined circadian rhythms in NK cell function at the gene, protein, and physiological levels under both basal and stimulated conditions. Finally, we sought to determine whether alterations in NK cell function due to repeated shifting of the LD cycle were related to an increased risk of tumor development, using an induction model of NK-sensitive adenocarcinoma cells (MADB106) (40, 41). MADB106 cancer cells typically metastasize in the lungs following i.v. injection, and the method is well established for exploring the role of NK cell function in tumor growth (41–43). We hypothesized that chronic shifting would suppress NK cell function in response to tumor cells by altering the coordinated circadian expression of key immune factors. We also postulated that these changes in NK cell function would promote tumor growth.

## Materials and Methods

### Animals and apparatus

Male Fischer (F344) rats were obtained from Charles River Laboratories (Wilmington, MA) at ~50 d of age. Upon arrival, animals were acclimated to the housing facility and maintained individually in running-wheel cages (wheel diameter: 34 cm; Mini Mitter, Bend, OR), placed within light- and sound-attenuating climate-controlled cubicles. Running-wheel activity was recorded by a computer and stored for later analysis of circadian parameters using the ClockLab software package (Actimetrics, Wilmette, IL). Animals were given free access to food and water for the duration of the experiment. All experimental procedures and animal treatment protocols were approved by Rutgers Animal Care and Facilities Committee and complied with National Institutes of Health policies.

### Procedures

Animals were maintained initially under a standard 12:12 LD cycle with lights on at 6:00 h and lights off at 18:00 h for a 1- to 2-wk acclimation period to stabilize running-wheel activity, and then assigned randomly to either remain under the standard LD cycle (controls) or to undergo the chronic shift-lag protocol ( $n = 90$  per lighting regimen used for various assays; see below). The shift-lag protocol consisted of 6-h LD phase advances repeated every 2 d for a total of 10 shifts (see Fig. 1A–C for a schematic representation of the control and shift-lag lighting regimens). For example, the first advance consisted of the lights coming on 6 h earlier than the previous day (lights on at 24:00 h), which resulted in a shortening of the dark period on the following day. Repeated phase advances were chosen owing to extensive prior data indicating that advances evoke greater circadian disruption than do phase delays. To determine whether the chronic shift-lag paradigm resulted in alterations of circadian expression of cytolytic factors, cytokines, and cytolytic capabilities in NK cells, a set of animals ( $n = 36$  per lighting regimen) were sacrificed by decapitation at six time points ( $n = 6$  per time point) 5–7 d into total darkness (DD), which corresponded to circadian times (CTs) 3, 7, 11, 19, and 23. To avoid the potential masking effects of light on these circadian rhythms, animals were sacrificed 5–7 d into DD following the last advance of the LD cycle. Actograms were used to determine CTs by defining CT12 as the onset of running-wheel activity. Immediately upon sacrifice, tissues were

collected and NK cells were separated from spleens for gene, protein, and cytotoxicity assays. To determine the effects of chronic shift-lag on NK cell cytotoxicity in response to tumor cells and the possible consequences on tumor growth, separate groups of animals ( $n = 30$  per lighting regimen) were inoculated with MADB106 cells at CT19 in DD. Half of these animals ( $n = 10$  per lighting regimen) were sacrificed 24 h after inoculation to assess immediate NK cell response to the presentation of tumor cells (40, 43). The remaining animals ( $n = 20$  per lighting regimen) were placed for 6–8 wk under the same standard LD cycle as during the acclimation period, and then sacrificed for lung extraction and inspection of tumor metastases.

### Enrichment of NK cells from spleen

Individual spleen tissue was processed as described previously (44). RBCs and granulocytes were removed from splenocyte suspensions by density centrifugation using Histopaque 1083 (Sigma-Aldrich). Splenocytes ( $\sim 10 \times 10^7$  cells per spleen) were extracted from the middle layer, washed with RPMI-1640 (Invitrogen), and resuspended in buffer (PBS, 0.5% BSA). Splenocytes were incubated with primary Abs conjugated to FITC (BD Biosciences), anti-CD6 (OX52; T cells), anti-CD45RA (OX-33; mature B cells), and anti-RT1B (OX-6; dendritic cells and macrophages), followed by secondary incubation with anti-FITC microbeads, according to the manufacturer's instructions (Miltenyi Biotec). NK cells were then enriched by magnetic separation (negative selection), using an AutoMACS Magnetic Separator (Miltenyi Biotec). The enriched fraction consistently yielded  $\sim 5 \times 10^6$  cells per spleen with a purity of  $\sim 80$ – $90\%$ , which was assessed by flow cytometry. CD3<sup>+</sup> and CD8<sup>+</sup> cells in the enriched fraction were consistently below 3%, and the remaining 10–20% of cells were believed to be a mixture of premature hematopoietic and endothelial cells. Following enrichment, NK cells were divided and lysed in appropriate buffers for RNA and protein analyses or were used for cytotoxicity assays.

### RNA extraction, RT-PCR, and real-time RT-PCR

Total RNA was isolated from  $\sim 2 \times 10^6$  NK cells per spleen, using the RNeasy Mini Kit (Qiagen). With the Superscript III First-Strand Synthesis SuperMix (Invitrogen) for RT-PCR, 100 ng total RNA was reverse transcribed and relative quantification of mRNA levels was performed by real-time RT-PCR (SYBR Green and TaqMan gene expression assays; Applied Biosystems), using an ABI Prism 7700 Sequence Detector. The following primer sequences were used. *Per2* F: 5'-GCAGCCTTTCGATTATTCTTC-3', R: 5'-GCTCCACGGGTTGATGAAG-3'; *Bmal1* F: 5'-TCCGATGACGAACCTGA-AACAC-3', R: 5'-CTCGGTCACATCCTACGACAA-3'; *perforin* F: 5'-GC-ATCGGTGCCAAGCCAGTG-3', R: 5'-GCCAGCGAGCCCCTGTCA-TCA-3', and specifically designed probes were used for granzyme-B and IFN- $\gamma$  (Applied Biosystems). Analyses were performed using the standard curve method with GAPDH (Applied Biosystems) as the normalizing endogenous control. Relative mRNA expression levels were calculated as the percentage of the maximum value over the 24-h period (45). The sample with the maximum ratio was used to compute percentages for all other samples by dividing each ratio by the maximum and converting to a percentage. Values at each CT represent a mean and SE of these values.

### Western blot

For protein analyses, enriched NK cells ( $\sim 2 \times 10^6$  cells per spleen) were lysed with a buffer containing protease and phosphatase inhibitors (25 mM Tris-HCl, pH 7.4; 150 mM NaCl; 1% Nonidet P-40; 1 mM EDTA; and 5% glycerol with Pierce Halt Protease Inhibitor). Protein (50  $\mu$ g) from total lysate was loaded and separated on a 4–20% SDS-PAGE and transferred overnight to polyvinylidene difluoride membranes. Membranes were incubated with primary Ab for 12 h at 4°C in blocking buffer, 5% w/v nonfat dry milk in TBS and 0.1% Tween 20. Primary Abs were used as follows: anti-BMAL1 rabbit polyclonal (1:500; Millipore), anti-PER2 mouse monoclonal (1:100; BD Biosciences), anti-perforin rabbit polyclonal (1:250), anti-granzyme B mouse monoclonal (1:200), and anti-IFN- $\gamma$  mouse monoclonal (1:250) Abs (Santa Cruz Biotechnology). Membranes were washed and incubated with peroxidase-conjugated secondary Abs (1:5000; Vector Laboratories) for 1 h at room temperature. Membranes were incubated with ECL Western blot chemiluminescence reagent (Pierce) and exposed to x-ray film, which was developed, and densitometry was performed using Image J analysis software (National Institutes of Health). Each protein was normalized to corresponding intensities for  $\beta$ -actin. Relative protein expression values were calculated as the percentage of the maximum value over the 24-h period (46). The sample with the maximum ratio was used to compute percentages for all other samples by dividing each ratio by the maximum and converting to a percentage. Values at each CT represent a mean and SE of these values.

### NK cell cytotoxicity

The cytotoxicity of enriched NK cells collected at each time point was determined by calcein AM assays using YAC-1 murine lymphoma cells (American Type Culture Collection) as target cells. YAC-1 cells were grown and maintained in RPMI 1640 without phenol red (Cellgro), containing 1% antibiotics (Sigma-Aldrich) and 10% FBS (Sigma-Aldrich). YAC-1 cells were washed and incubated with 5 mM calcein AM (Sigma-Aldrich) in serum-free RPMI-1640 for 10 min at 37°C. Labeled YAC-1 cells were washed and plated into U-bottom 96-well plates (Falcon) at a concentration of  $5 \times 10^4$  cells per well. NK cells were added at various E:T ratios in triplicates. YAC-1 cells in RPMI alone were to determine spontaneous calcein AM release, whereas maximal release was achieved by lysing target cells in buffer (0.1% Triton X-100). NK cells collected over the circadian timescale were preincubated for 12 h with IL-2 at 37°C (100 ng/ml; R&D Systems) prior to 4-h incubation with YAC-1 target cells. IL-2 was used to increase assay efficiency owing to the comparatively selective nature of this cytokine to increase cytotoxicity in NK cells (47). NK cells collected from animals inoculated with MADB106 cells at CT19 were incubated at 37°C for 4 h with YAC-1 cells in the absence of IL-2. All assays were analyzed using a fluorescence plate reader (Tecan). The percent cytotoxicity for each sample was calculated as follows: % cytotoxicity = [(experimental well – spontaneous well)/(max lysed well – spontaneous well)]  $\times$  100. The percentages at each E:T ratio were converted to LUs per  $10^7$  effector cells and based on 20% specific cell lysis (48).

### Tumor cell inoculation and metastases

Following control and chronic shift-lag protocols, animals were placed under DD for 5–7 d and inoculated at CT19 with MADB106 tumor cells ( $1 \times 10^5$  cells/0.2 ml/rat) into the jugular vein, under isoflurane anesthesia (Henry Schein). As noted above, CT19 was determined by wheel-running data for each animal. This time was chosen based on previous studies and our pilot data showing a maximal response of NK cells during the dark phase of the LD cycle (45, 46). Isoflurane was used to minimize the amount of time the animal was anesthetized. Small incision sites (~1 cm) were closed with a surgical clip, and rats were placed back into their home cage. Surgical and inoculation procedures took ~5–7 min to complete per animal. Animals were sacrificed 6–8 wk after tumor cell inoculation, and each animal was inspected for the presence of visible tumors. As expected, tumors were located primarily in the lungs. Lungs were collected and placed for 24 h in Bouin's solution (72% saturated picric acid solution, 23% formaldehyde 37% solution, and 5% glacial acetic acid). Lungs were washed with ethanol, and two individuals separately counted surface tumor metastases.

### Plasma analysis of corticosterone

Blood samples (~80–100  $\mu$ l) were collected at CT7 and CT19 in 15  $\mu$ l EDTA and centrifuged for plasma. Plasma was analyzed for corticosterone levels by a competitive ELISA (IBL International) according to the manufacturer's recommendations. All samples from both groups were run on a single 96-well plate to limit variability. Standard curves were used to determine corticosterone levels.

### Data analysis

Two-way ANOVA with Bonferroni post hoc tests was used to determine differences between control and shift-lag groups at each circadian phase. The gene and protein expression data were tested for significant circadian rhythmicity, using CircWave v. 1.4 software (21, 49) to determine the best-fitting linear harmonic regression with an assumed period of 24 h and with  $\alpha$  set at 0.05. The center of gravity of each best-fitting waveform, which comprises both sine and cosine components, in CircWave was used as the circadian acrophase, and the associated estimation error was used as the SD. In higher order waveforms, the center of gravity is a reliable estimator of phase of peak expression (50, 51). One-way ANOVA with Newman–Keuls multiple comparison tests was used to compare acrophases between groups for each gene or protein. Student *t* tests were used to compare group means for circadian acrophases of NK cell activity and protein and gene expression, tumor frequency data, and circadian periods. The Mann–Whitney *U* test was used to compare tumor frequencies between different defined chronotypes of the chronic shift-lag animals. Circadian period and robustness of running-wheel activity rhythms were derived using Lomb–Scargle periodograms during and following the control and shift-lag protocols (ClockLab).

## Results

### Circadian wheel-running activity rhythms

Control animals under the standard LD cycle displayed typical circadian entrainment patterns (Fig. 1A) with an approximate pe-


riod of 24 h (Fig. 1D). Under the chronic shift-lag protocol, rats exhibited different patterns of locomotor activity rhythms. For most shift-lag animals, periodogram analysis revealed two distinct peaks, one corresponding to the mean period of the shift-lag schedule (i.e., ~21 h) and one corresponding to the typical, longer-than-24 h free-running period normally shown by rats under free-running conditions (Fig. 1E, 1F). However, individual variation was considerable among shift-lag animals in the relative prominence of these peaks and in a clear correspondence between the periodogram results and the actigraphic record within individuals (Fig. 1B, 1C). In DD following the shifting paradigm, rhythm amplitude was significantly reduced, with no observable change in period (Fig. 2A, 2B;  $p < 0.0001$ ). Thus, the shift-lag protocol induced considerable circadian disruption and also resulted in light exposure during the animals' subjective night.

### Chronic shift-lag alters the circadian expression of clock genes in enriched NK cells

To determine whether chronic shifting affected the molecular clock in peripheral immune cells, we examined gene and protein expression of two canonical clock genes, *Per2* and *Bmal1* (52), from enriched NK cells collected during DD at 4-h intervals across the circadian cycle. In control animals, both *Per2* and *Bmal1* gene and protein expression displayed robust circadian rhythmicity (Fig. 3; all  $p < 0.01$ ). Acrophase estimates (estimate  $\pm$  SD) derived from fitted curves indicated *Per2* and *Bmal1* expression peaks were out of phase with one another at both the gene (Fig. 3A, 3C) and the protein (Fig. 3B, 3D) levels (Table I), as expected from previous studies (45). In addition, as hypothesized, chronic shift-lag altered the circadian expression of both clock genes and proteins; nevertheless, these displayed significant rhythmicity in shift-lag animals (Fig. 3; all  $p < 0.01$ ). Comparison of acrophases for *Per2* and *Bmal1* revealed that expression rhythms were shifted in NK cells from shift-lagged animals compared with controls (Table I; Fig. 3A, 3C;  $p < 0.05$ ), whereas *BMAL1*, but not *PER2*, phase of peak expression was affected by shift-lag (Table I; Fig. 3B, 3D). Moreover, two-way ANOVAs indicated significant group by time interactions ( $p < 0.05$ ) for *Per2* and *Bmal1* gene and protein expression rhythms, which were further explored using Bonferroni post hoc tests to indicate significant changes in levels of expression at each CT. Levels of expression were significantly altered at particular CTs for *Per2* and *Bmal1* (Fig. 3), further indicating chronic shift-lag affected the rhythmic expression of core clock genes, with the most profound effects on *BMAL1* expression.

### Chronic shift-lag alters the circadian expression of the cytokine IFN- $\gamma$ and cytolytic factors perforin and granzyme B in enriched NK cells

Alterations in circadian expression of core components of the molecular clock were paralleled by changes in expression rhythms of cytokines and cytolytic factors in NK cells. Consistent with previous results (37, 38), NK cells from control animals displayed robust circadian rhythms of gene and protein expression of perforin, granzyme B, and IFN- $\gamma$  (Fig. 4; assumed period of 24 h, all  $p < 0.01$ ), with expression peaks during the active phase (Table I). In line with our hypothesis, chronic shift-lag altered the rhythmic gene and protein expression of perforin, granzyme B, and IFN- $\gamma$  (Fig. 4). Chronic shift-lag completely abolished the gene and protein rhythms of perforin in NK cells, as indicated by a lack of significant curve fitting and decreased gene expression across the entire circadian cycle (Fig. 4A, 4B; compared with controls, all CTs,  $p < 0.05$ ). Although gene expression rhythms remained intact for granzyme B and IFN- $\gamma$  in animals undergoing shift-lag ( $p < 0.01$ ), these gene rhythms were shifted to earlier in the circadian cycle, as indicated



**FIGURE 1.** Effects of chronic shift-lag on circadian wheel-running activity rhythms. Circadian wheel-running activity rhythms depicted as standard raster-style double-plotted actograms from several representative animals from control (**A**) and shifted (**B**, **C**) groups. Repeated shifts of the LD cycle resulted in different behavioral patterns of re-entrainment among animals. Shift-lagged animals tended to essentially “ignore” changing environmental light cues (**B**) or re-entrain to them (**C**). LS periodogram analysis for control (**D**) and shifted (**E**) animals of rhythm period and amplitude during standard or shifted LD cycles revealed that shifted animals tend to remain rhythmic with a significantly longer period ( $*p < 0.0001$ ). Large secondary peaks are evident in the shifted group (arrow), which, with further analysis, revealed particular animals displaying shorter periods. Robust periods shorter in length were also common in the shifted group, which is represented by an actogram (**C**) and periodogram (**F**) of an individual animal during the shifting paradigm. No differences were found in rhythm amplitude during standard or shifting paradigms. Shaded regions represent the dark phase of the LD cycle.

by changes in acrophase (Table I) and changes in expression levels at particular phases (Fig. 4C, 4E). The protein expression of granzyme B and IFN- $\gamma$  did not show a similar shift in peak expression (Table I; Fig. 4D, 4F). In general, alterations in rhythms of these factors due to shift-lag resulted in reductions of gene and protein levels near or at phases of peak expression in control animals (Fig. 4;  $p < 0.05$ ). Therefore, chronic shift-lag appeared to differentially alter gene and protein expression of cytokines and cytolytic factors by either shifting the circadian rhythm to earlier in the cycle or attenuating levels at phases of peak expression.


#### *Chronic shift-lag alters the circadian rhythmicity of NK cell cytolytic function*

In an effort to investigate whether changes in circadian rhythms of cytokines and cytolytic factors due to chronic shift-lag had consequences for NK cell function, we completed a series of NK cell cytotoxicity assays across the circadian cycle. The cytotoxic function of NK cells displays a robust circadian rhythm with a peak during the subjective night (Table I; Fig. 5A;  $p < 0.001$ ), which is


in line with our previous results (46). In shifted animals, NK cell cytotoxicity remained rhythmic, with a shift in the rhythm by  $\sim 16$  h (Table I; Fig. 5A;  $p < 0.05$ ). Importantly, it is evident that peaks of NK cell cytotoxicity observed in controls animals were dramatically reduced in the shifted group (Fig. 5A;  $p < 0.05$ ). From these data, it is suggested that coordinated circadian peaks of expression of perforin, granzyme B, and IFN- $\gamma$  are important for NK cell-mediated killing.

To determine whether chronic shift-lag altered NK cell function *in vivo*, enriched NK cells were collected from animals injected with MADB106 tumor cells. On the premise that circadian activity of NK cell function is critical for immune response to cancer, tumor cells were injected at CT19—the phase of maximal cytolytic activity (Table I; Fig. 5A). As expected, NK cells from shifted animals collected 24 h following the *i.v.* introduction of tumor cells displayed significantly reduced cytolytic activity (Fig. 5B;  $p < 0.01$ ). From these data, we interpret that shift-lag significantly altered the circadian rhythm of NK cell cytolytic function and the NK cells’ response to tumor challenge.

**FIGURE 2.** Circadian wheel-running activity rhythm amplitudes are decreased following chronic shift-lag. LS periodogram analysis was restricted to the initial 5 d in DD for control (**A**) and shifted (**B**) groups. No differences for period were demonstrated between groups in DD. Rhythm amplitude levels (dotted lines) were significantly decreased in DD following the chronic shifting paradigm ( $*p < 0.0001$ ). Shifted animals remained comparatively rhythmic with a dampening of circadian rhythm.



**FIGURE 3.** Chronic shift-lag alters the circadian expression of clock genes and proteins in enriched NK cells. Expression levels of *Bmal1* and *Per2* genes (**A, C**) and associated proteins (**B, D**) in NK cells separated from animals at different time points in DD following either control or shifted paradigms. Representative immunoblots are shown above respective protein plots across all CTs. Densitometric quantification of protein levels was done by Image J software. Sine wave fits using linear harmonic regression with an assumed period of 24 h for control and shifted groups (solid and dotted lines, respectively) are superimposed with group means  $\pm$  SEMs for each CT. All curve fits are significant ( $p < 0.01$ ). \* $p < 0.05$ , significant difference between groups at CT.



*Decreased innate immune function in chronically shifted animals is associated with increases in lung tumor frequency and prevalence*

Control and shift-lag animals injected at CT19 with MADB106 tumor cells were left for 6–8 wk to determine tumor frequency or prevalence. In line with our hypothesis, chronic shift-lag promoted tumor growth, as indicated by increased frequency and prevalence of tumors in the lungs (Fig. 6A, 6B;  $p < 0.001$ ). In addition, tumor load was compared between animals displaying two different chronotypes in response to the shifting protocol—animals that exhibited either free-running activity rhythms or re-entrainment (Fig. 1B, 1C, respectively). The shifted group that displayed repeated re-entrainment tended to have more tumors relative to the group that did not successfully synchronize to the shift-lag schedule, but this difference did not reach significance (Fig. 6C;  $p < 0.07$ ). Representative lungs from control and shifted (Fig. 6D, rows 1 and 2, respectively) animals are pictured. Two animals

from the chronic shift-lag group were removed from the analysis because tumor growth was so invasive to the lungs and surrounding organs that each required sacrificing 3 wk following injection (not included in the analysis). No control animals required sacrificing prior to the planned 6–8 wk.

*Suppressed NK cell function in animals undergoing chronic shift-lag may not be due to increases in plasma corticosterone*

Other studies have shown that repeated shifting of the LD cycle either shifts the rhythm or transiently increases concentration of plasma corticosterone (15, 53). To determine whether shifting increased plasma corticosterone, levels were measured at two time points, CT7 and CT19, at the time of sacrifice (5–7 d into DD;  $n = 4$  per time point). Corticosterone levels did not differ between groups at either time point (Fig. 7). At CT19, the time of maximal NK cytolytic activity in controls, corticosterone levels were not increased in shifted animals, suggesting that attenuation of NK function may not be due to activation of the stress axis.


Table I. Chronic shift-lag alters the acrophases of gene, protein, and cytolytic activity rhythms in NK cells

	Gene		Protein	
	Control	Shifted	Control	Shifted
Per2	9.54 $\pm$ 3.26*	17.71 $\pm$ 3.30**	14.2 $\pm$ 3.49*	15.28 $\pm$ 3.25
Bmal1	20.88 $\pm$ 2.21	13.03 $\pm$ 2.77**	19.05 $\pm$ 2.54	9.38 $\pm$ 3.32**
Perf	22.24 $\pm$ 3.23	—	18.85 $\pm$ 3.18	—
Gran B	20.66 $\pm$ 1.76	13.66 $\pm$ 3.46**	18.90 $\pm$ 3.10	19.32 $\pm$ 2.96
IFN- $\gamma$	18.97 $\pm$ 2.46	13.95 $\pm$ 3.00**	20.90 $\pm$ 3.32	20.93 $\pm$ 3.12
	Cytolytic Activity			
	Control		Shifted	
NK cells	17.35 $\pm$ 3.04		3.78 $\pm$ 3.25**	

Acrophases were determined from curves fitted to gene and protein expression and NK cell activity levels over the circadian cycle. Acrophases ( $\pm$ SD) represent estimations of the circadian phase corresponding to the peak of the rhythm. One-way ANOVA with Newman-Keuls multiple comparison tests were used to compare acrophases between control and shifted groups ( $n = 6$  per lighting regimen) by gene and protein, or, in the case of NK cell activity ( $n = 4$  per lighting regimen), a Student  $t$  test was used.

\* $p < 0.05$ , significant differences between groups, \*\* $p < 0.05$ , significant differences between Per2 and Bmal1.


**FIGURE 4.** Chronic shift-lag alters the circadian expression of cytokines and cytolytic factors in enriched NK cells. Expression levels of genes (*left*) and proteins (*right*) for cytolytic factors perforin (**A, B**) and granzyme B (**C, D**) and for cytokine IFN- $\gamma$  (**E, F**) in NK cells separated from animals at different time points in DD following either control or shifted paradigms. Representative immunoblots are shown above respective protein plots across all CTs. Densitometric quantification of protein levels was done by Image J software. Sine wave fits using linear harmonic regression with an assumed period of 24 h for control and shifted groups (solid and dotted lines, respectively) are superimposed with group means  $\pm$  SEMs for each CT. Straight line (i.e., perforin) denotes no significant curve fit, and thus a complete lack of rhythm. All curve fits are significant ( $p < 0.01$ ). \* $p < 0.05$ , significant difference between groups at CT.



## Discussion


The circadian system regulates numerous cellular and physiological processes, including immune function (34). At the top of

the circadian hierarchy lies the SCN, which coordinates timing of peripheral clocks by hormonal and autonomic nervous system signals (54). For example, immune tissues receive rhythmic sympathetic input that modulates clock gene expression and other circadian functions within immunomodulatory cells, including splenic NK cells (37, 55, 56). NK cell function seems to be pervasively regulated by clock mechanisms, as clock genes, cytokines, and cytolytic factors all oscillate in a circadian fashion (38, 45). The present study demonstrates that circadian coordination of the immune response may influence the degree to which the innate immune system initially reacts to circulating tumor cells. During a state of chronic circadian desynchrony, the cytolytic activity of NK cells is almost completely suppressed, quite possibly by disruption of the coordinated expression of multiple cytolytic factors. We postulate that this mechanism is responsible for the increased incidence of lung tumors in animals undergoing chronic shift-lag.



**FIGURE 5.** Cytotoxicity of NK cells is suppressed in animals undergoing chronic shift-lag. NK cell cytotoxicity expresses a robust circadian rhythm, which is significantly suppressed during peak times by chronic shift-lag (**A**). Under immune-stimulated conditions, NK cell cytotoxicity is decreased in animals undergoing chronic shift-lag (**B**). Sine wave fits using linear harmonic regression with an assumed period of 24 h for control and shifted groups (solid and dotted lines, respectively) are superimposed with group means  $\pm$  SEMs for each CT. All curve fits are significant ( $p < 0.01$ ). \* $p < 0.05$ , significant difference between groups at CT; \*\* $p < 0.01$ , significant difference between groups.

A number of epidemiological studies have linked disruptions in circadian rhythms to cancer susceptibility and clinical course, although the genetic and physiological mechanisms underlying these findings are poorly understood. Much attention has been focused on the role of the molecular clock in cell proliferation, DNA damage and repair, and apoptotic pathways within the developing tumor cells (16, 57–59). In contrast, the current study shows that the tumorigenic effects of circadian disruption depend on tumor surveillance mechanisms of NK cells. Specifically,



**FIGURE 6.** Chronic shift-lag increases tumor frequency and incidence in the lungs. Lung tumor frequency (**A**) and percentage of animals with tumors (**B**) are increased in shifted animals. Animals displaying re-entrainment during the repeated shifting schedule tended to have an increased tumor load, compared with shifted animals displaying free-running activity rhythms (**C**;  $p = 0.07$ ). Pictures of lungs with multiple tumor loci (circled) from representative animals in control (**D**, row 1) and shifted (**D**, row 2) groups. \*\*\* $p < 0.0001$ , significant difference between groups.

chronic shift-lag altered the expression of clock genes and the immunomodulators IFN- $\gamma$ , perforin, and granzyme B in enriched NK cells at both the gene and the protein levels. Thus, the functional activity of NK cells was significantly decreased at normal phases of peak expression in shift-lag compared with control animals. Further, gene expression rhythms of granzyme B and IFN- $\gamma$  displayed a shift by  $\sim 4$  h to earlier in the circadian cycle, but similar shifts were not observed at the protein level. Perforin expression appeared the most affected by chronic shift-lag, as gene and protein rhythms were completely abolished in NK cells. It is plausible that these reductions of perforin hamper NK cell-mediated apoptosis and lysis of tumor cells in shifted animals (60–62), especially because granzyme B and IFN- $\gamma$  were also reduced, which has been associated with increased tumor susceptibility (31).


Circadian coordination of NK cell function appears to be important for generating an efficient immune response. Consistent with our previous findings, circadian rhythms of perforin, granzyme B, and IFN- $\gamma$  expression peak at similar phases during the subjective night, corresponding to the circadian peak of NK cell cytotoxicity (45, 46). In an effort to determine whether circadian coordination of NK cell function may have functional significance, we injected control and shifted animals with tumor cells at CT19 according to the hypothesis that chronic shift-lag would cause impairment of NK cell function, ultimately producing a shift in the rhythms of cytotoxicity. Somewhat surprisingly, however, chronic shift-lag dramatically attenuated NK cell function across the entire circadian cycle, suggesting that similar tumor frequencies might have been observed in shifted animals regardless of phase of tumor cell injection. The apparent discoordination of

cytokines, cytolytic factors, and cytotoxicity, along with the attenuation of cytotoxicity following an *in vivo* tumor challenge, suggests that circadian disruption compromises the ability of NK cells to kill tumor cells. In our model, this compromise could lead to the promotion of tumor growth by increasing the probability of tumor cell retention following inoculation.

To assess circadian disruption at the whole-animal level, we monitored circadian activity rhythms throughout the experimental protocols. Because rhythmic locomotor activity is a reliable indicator of circadian pacemaker functioning, changes in amplitude and/or period due to chronic shift-lag can be inferred to reflect SCN activity. Inspection of actograms from our shifted animals yielded two different chronotypes. Although some shifted animals exhibited an apparent free-running period, essentially ignoring the changing light onsets, others displayed short circadian periods as a consequence of partial synchronization to the advancing shift-lag schedule. Using a similar shifting protocol with hamsters, the former has been observed (53). Nevertheless, despite shifting-induced alterations in rhythm amplitude and period, our animals remained completely rhythmic during the experiment and thus suggest the SCN maintains coherent rhythmicity even during rapidly changing environmental cues (53). Studies examining the effects of photic phase shifts on SCN clock gene expression have also demonstrated the SCN maintains rhythmicity with differing periods of re-entrainment, depending on frequency and direction of the shifts (63).

Furthermore, examination of circadian activity rhythms during the exposure to the shift-lag schedule revealed that repeated re-entrainment may have led to increased tumorigenesis relative to animals expressing stable free-running rhythms under the same conditions. Because re-entrainment is associated with internal circadian desynchrony at both the molecular and the physiological levels (64, 65), these preliminary observations support the hypothesis that dissociation among multiple circadian clocks may indeed underlie the pathophysiological effects of shift-lag exposure.

Nevertheless, although rhythmicity is maintained in the SCN, major SCN outputs that relay temporal information to other brain areas and peripheral tissues are compromised under shift-lag conditions (63, 64). Clocks in peripheral tissues re-entrain at a slower rate, compared with the SCN, following a single advance or delay, including those in the spleen (63, 65). Sympathetic input to the spleen may be a major output pathway by which the SCN coordinates NK cell function (37). It is interesting to speculate that



**FIGURE 7.** Plasma corticosterone levels in control and shifted animals. Corticosterone was measured at two circadian phases, CT7 and CT19, at the time of sacrifice 5–7 d into DD. No differences were observed between groups at either time point. Suppressed NK cell function may not be due to increases in corticosterone levels.



chronic shift-lag impedes temporal coordination of the immune system by attenuating SCN-derived output signals to “preautonomic” neurons (66). Indeed, homeostasis of the sympathetic nervous system is disrupted in chronically shifted animals (17).

The molecular clock in NK cells was altered by chronic shift-lag, which was evident by advances in the rhythms and reductions of *Bmal1* gene and protein expression, as well as a delay in the expression rhythms of *Per2*. *Bmal1* rhythms were slightly attenuated but remained rhythmic. The delay of the *Per2* gene was much more dramatic (~6 h) than the pattern observed in the protein rhythm (~1 h). Although shifting-induced reductions in NK cell function were correlated with changes in clock gene expression, these data can only suggest the involvement of molecular clocks in NK cells in regulating cytolytic abilities. Similarly, in macrophages, disruptions to clock genes due to repeated shifting has been associated with dysregulation of inflammatory responses (20). To determine the extent to which molecular clocks within NK cells regulate cell-specific functions, further studies are necessary to tease apart the interactions between clock and immune-related pathways. At the molecular level, a potential intersection could be clock modulation of NF- $\kappa$ B, a key transcription factor involved in cellular immune response (67). When activated, NF- $\kappa$ B can promote gene transcription of IFN- $\gamma$  (68), perforin (69), and granzyme B (70) by direct binding to enhancer elements.

Other factors could be involved in mediating the effects of chronic shift-lag on NK cell function beyond involvement of circadian mechanisms. Circulating glucocorticoids and melatonin oscillate in antiphase, peaking during the early and late circadian cycle, respectively, and are known modulators of NK cell function (71, 72). However, several points of evidence suggest glucocorticoids and melatonin may not necessarily be critical for maintaining immune rhythms. In macrophages and NK cells, functional rhythms persist despite the absence of glucocorticoid rhythms (73, 74) or lack of melatonin production (73, 75). Furthermore, consistent with a previous result (20), our study suggests dysregulation of NK cell function induced by chronic shift-lag is not due to increased circulating corticosterone, especially during times of peak functioning. Corticosterone increases induced by chronic shifting paradigms may be transient and play a role in resynchronization of the circadian system (49, 53). Nevertheless, at the time points of NK cell collection, corticosterone was not increased in our shifted animals.

In conclusion, we have demonstrated that chronic states of circadian desynchrony severely attenuated NK cell function, possibly by altering the circadian rhythms of cytokines, cytolytic factors, and cytotoxicity. In addition, suppressed NK cell function was observed during an in vivo immune-stimulated condition in shifted animals, suggesting circadian disruption has substantial effects on the tumor response of NK cells. Importantly, loss of rhythms in NK cell cytolytic activity was associated with an increased frequency of tumors. More detailed studies are required to further understand the integration of physiological and molecular changes in immune function caused by disruption to the circadian system. For example, whether the chronic shift-lag model used in this study alters noradrenergic signaling of the sympathetic nervous system to impact NK cell function has yet to be determined (17). Molecular studies, such as large-scale gene expression arrays in primary and secondary immune tissues, of wild-type and clock mutant mice may also provide a wealth of information regarding the molecular links between circadian and immune systems. Understanding these intersections could be invaluable to developing novel therapeutics to restore circadian functioning in immune cells, which could aid in preventing disease progression (59, 76, 77).

## Disclosures

The authors have no financial conflicts of interest.

## References

- Lahti, T. A., T. Partonen, P. Kyrrönen, T. Kauppinen, and E. Pukkala. 2008. Night-time work predisposes to non-Hodgkin lymphoma. *Int. J. Cancer* 123: 2148–2151.
- Davis, S., D. K. Mirick, and R. G. Stevens. 2001. Night shift work, light at night, and risk of breast cancer. *J. Natl. Cancer Inst.* 93: 1557–1562.
- Hansen, J. 2001. Light at night, shiftwork, and breast cancer risk. *J. Natl. Cancer Inst.* 93: 1513–1515.
- Schernhammer, E. S., F. Laden, F. E. Speizer, W. C. Willett, D. J. Hunter, I. Kawachi, and G. A. Colditz. 2001. Rotating night shifts and risk of breast cancer in women participating in the nurses' health study. *J. Natl. Cancer Inst.* 93: 1563–1568.
- Viswanathan, A. N., and E. S. Schernhammer. 2009. Circulating melatonin and the risk of breast and endometrial cancer in women. *Cancer Lett.* 281: 1–7.
- Conlon, M., N. Lightfoot, and N. Kreiger. 2007. Rotating shift work and risk of prostate cancer. *Epidemiology* 18: 182–183.
- Kubo, T., K. Ozasa, K. Mikami, K. Wakai, Y. Fujino, Y. Watanabe, T. Miki, M. Nakao, K. Hayashi, K. Suzuki, et al. 2006. Prospective cohort study of the risk of prostate cancer among rotating-shift workers: findings from the Japan collaborative cohort study. *Am. J. Epidemiol.* 164: 549–555.
- Kloog, I., A. Haim, R. G. Stevens, and B. A. Portnov. 2009. Global co-distribution of light at night (LAN) and cancers of prostate, colon, and lung in men. *Chronobiol. Int.* 26: 108–125.
- Schernhammer, E. S., F. Laden, F. E. Speizer, W. C. Willett, D. J. Hunter, I. Kawachi, C. S. Fuchs, and G. A. Colditz. 2003. Night-shift work and risk of colorectal cancer in the nurses' health study. *J. Natl. Cancer Inst.* 95: 825–828.
- Mormont, M. C., J. Waterhouse, P. Bleuzen, S. Giacchetti, A. Jami, A. Bogdan, J. Lellouch, J. L. Misset, Y. Touitou, and F. Lévi. 2000. Marked 24-h rest/activity rhythms are associated with better quality of life, better response, and longer survival in patients with metastatic colorectal cancer and good performance status. *Clin. Cancer Res.* 6: 3038–3045.
- Sephton, S. E., R. M. Sapolsky, H. C. Kraemer, and D. Spiegel. 2000. Diurnal cortisol rhythm as a predictor of breast cancer survival. *J. Natl. Cancer Inst.* 92: 994–1000.
- Chaput, J. P., M. Brunet, and A. Tremblay. 2006. Relationship between short sleeping hours and childhood overweight/obesity: results from the “Québec en Forme” Project. *Int. J. Obes. (Lond.)* 30: 1080–1085.
- Morikawa, Y., H. Nakagawa, K. Miura, Y. Soyama, M. Ishizaki, T. Kido, Y. Naruse, Y. Suwazono, and K. Nogawa. 2005. Shift work and the risk of diabetes mellitus among Japanese male factory workers. *Scand. J. Work Environ. Health* 31: 179–183.
- Tüchsen, F., H. Hamnerz, and H. Burr. 2006. A 12 year prospective study of circulatory disease among Danish shift workers. *Occup. Environ. Med.* 63: 451–455.
- Filipksi, E., F. Delaunay, V. M. King, M. W. Wu, B. Claustrat, A. Gréchez-Cassiau, C. Guettier, M. H. Hastings, and L. Francis. 2004. Effects of chronic jet lag on tumor progression in mice. *Cancer Res.* 64: 7879–7885.
- Filipksi, E., X. M. Li, and F. Lévi. 2006. Disruption of circadian coordination and malignant growth. *Cancer Causes Control* 17: 509–514.
- Lee, S., L. A. Donehower, A. J. Herron, D. D. Moore, and L. Fu. 2010. Disrupting circadian homeostasis of sympathetic signaling promotes tumor development in mice. *PLoS ONE* 5: e10995.
- Pendergast, J. S., M. Yeom, B. A. Reyes, Y. Ohmiya, and S. Yamazaki. 2010. Disconnected circadian and cell cycles in a tumor-driven cell line. *Commun. Integr. Biol.* 3: 536–539.
- Yu, E. A., and D. R. Weaver. 2011. Disrupting the circadian clock: gene-specific effects on aging, cancer, and other phenotypes. *Aging (Albany NY)* 3: 479–493.
- Castanon-Cervantes, O., M. Wu, J. C. Ehlen, K. Paul, K. L. Gamble, R. L. Johnson, R. C. Besing, M. Menaker, A. T. Gewirtz, and A. J. Davidson. 2010. Dysregulation of inflammatory responses by chronic circadian disruption. *J. Immunol.* 185: 5796–5805.
- Keller, M., J. Mazuch, U. Abraham, G. D. Eom, E. D. Herzog, H. D. Volk, A. Kramer, and B. Maier. 2009. A circadian clock in macrophages controls inflammatory immune responses. *Proc. Natl. Acad. Sci. USA* 106: 21407–21412.
- Zamai, L., C. Ponti, P. Mirandola, G. Gobbi, S. Papa, L. Galeotti, L. Cocco, and M. Vitale. 2007. NK cells and cancer. *J. Immunol.* 178: 4011–4016.
- Dunn, G. P., L. J. Old, and R. D. Schreiber. 2004. The immunobiology of cancer immunosurveillance and immunoediting. *Immunity* 21: 137–148.
- Smyth, M. J., J. Swann, E. Cretney, N. Zerafa, W. M. Yokoyama, and Y. Hayakawa. 2005. NKG2D function protects the host from tumor initiation. *J. Exp. Med.* 202: 583–588.
- Street, S. E., Y. Hayakawa, Y. Zhan, A. M. Lew, D. MacGregor, A. M. Jamieson, A. Diefenbach, H. Yagita, D. I. Godfrey, and M. J. Smyth. 2004. Innate immune surveillance of spontaneous B cell lymphomas by natural killer cells and gamma-delta T cells. *J. Exp. Med.* 199: 879–884.
- Imai, K., S. Matsuyama, S. Miyake, K. Suga, and K. Nakachi. 2000. Natural cytotoxic activity of peripheral-blood lymphocytes and cancer incidence: an 11-year follow-up study of a general population. *Lancet* 356: 1795–1799.
- Konjevic, G., S. Radenkovic, T. Srdic, V. Jurisic, Lj. Stamatovic, and M. Milovic. 2011. Association of decreased NK cell activity and IFN $\gamma$  expression with pSTAT dysregulation in breast cancer patients. *J. BUON* 16: 219–226.

28. Cullen, S. P., M. Brunet, and S. J. Martin. 2010. Granzymes in cancer and immunity. *Cell Death Differ.* 17: 616–623.
29. Vivier, E., E. Tomasello, M. Baratin, T. Walzer, and S. Ugolini. 2008. Functions of natural killer cells. *Nat. Immunol.* 9: 503–510.
30. Cifone, M. G., S. D'Alò, R. Parroni, D. Millimaggi, L. Biondi, S. Martinotti, and A. Santoni. 1999. Interleukin-2-activated natural killer cells express inducible nitric oxide synthase that contributes to cytotoxic function and interferon-gamma production. *Blood* 93: 3876–3884.
31. Lee, S. H., T. Miyagi, and C. A. Biron. 2007. Keeping NK cells in highly regulated antiviral warfare. *Trends Immunol.* 28: 252–259.
32. Shankaran, V., H. Ikeda, A. T. Bruce, J. M. White, P. E. Swanson, L. J. Old, and R. D. Schreiber. 2001. IFN-gamma and lymphocytes prevent primary tumour development and shape tumour immunogenicity. *Nature* 410: 1107–1111.
33. Street, S. E., E. Cretney, and M. J. Smyth. 2001. Perforin and interferon-gamma activities independently control tumor initiation, growth, and metastasis. *Blood* 97: 192–197.
34. Logan, R. W., and D. K. Sarkar. 2012. Circadian nature of immune function. *Mol. Cell. Endocrinol.* 349: 82–90.
35. Albrecht, U., and H. Oster. 2001. The circadian clock and behavior. *Behav. Brain Res.* 125: 89–91.
36. Silver, R., J. LeSauter, P. A. Tresco, and M. N. Lehman. 1996. A diffusible coupling signal from the transplanted suprachiasmatic nucleus controlling circadian locomotor rhythms. *Nature* 382: 810–813.
37. Logan, R. W., A. Arjona, and D. K. Sarkar. 2011. Role of sympathetic nervous system in the entrainment of circadian natural-killer cell function. *Brain Behav. Immun.* 25: 101–109.
38. Arjona, A., and D. K. Sarkar. 2006. Evidence supporting a circadian control of natural killer cell function. *Brain Behav. Immun.* 20: 469–476.
39. Arjona, A., and D. K. Sarkar. 2008. Are circadian rhythms the code of hypothalamic-immune communication? Insights from natural killer cells. *Neurochem. Res.* 33: 708–718.
40. Ben-Eliyahu, S., G. G. Page, R. Yirmiya, and A. N. Taylor. 1996. Acute alcohol intoxication suppresses natural killer cell activity and promotes tumor metastasis. *Nat. Med.* 2: 457–460.
41. Melamed, R., S. Bar-Yosef, G. Shakhar, K. Shakhar, and S. Ben-Eliyahu. 2003. Suppression of natural killer cell activity and promotion of tumor metastasis by ketamine, thiopental, and halothane, but not by propofol: mediating mechanisms and prophylactic measures. *Anesth. Analg.* 97: 1331–1339.
42. Page, G. G., and S. Ben-Eliyahu. 1999. A role for NK cells in greater susceptibility of young rats to metastatic formation. *Dev. Comp. Immunol.* 23: 87–96.
43. Sarkar, D. K., C. Zhang, S. Murugan, M. Dokur, N. I. Boyadjieva, M. Ortigueira, K. R. Reuhl, and S. Mojtehdzadeh. 2011. Transplantation of  $\beta$ -endorphin neurons into the hypothalamus promotes immune function and restricts the growth and metastasis of mammary carcinoma. *Cancer Res.* 71: 6514–6523.
44. Boyadjieva, N., M. Dokur, J. P. Advis, G. G. Meadows, and D. K. Sarkar. 2001. Chronic ethanol inhibits NK cell cytolytic activity: role of opioid peptide beta-endorphin. *J. Immunol.* 167: 5645–5652.
45. Arjona, A., and D. K. Sarkar. 2005. Circadian oscillations of clock genes, cytolytic factors, and cytokines in rat NK cells. *J. Immunol.* 174: 7618–7624.
46. Arjona, A., N. Boyadjieva, and D. K. Sarkar. 2004. Circadian rhythms of granzyme B, perforin, IFN-gamma, and NK cell cytolytic activity in the spleen: effects of chronic ethanol. *J. Immunol.* 172: 2811–2817.
47. Wang, K. S., J. Ritz, and D. A. Frank. 1999. IL-2 induces STAT4 activation in primary NK cells and NK cell lines, but not in T cells. *J. Immunol.* 162: 299–304.
48. Bryant, J., R. Day, T. L. Whiteside, and R. B. Herberman. 1992. Calculation of lytic units for the expression of cell-mediated cytotoxicity. *J. Immunol. Methods* 146: 91–103.
49. Kiessling, S., G. Eichele, and H. Oster. 2010. Adrenal glucocorticoids have a key role in circadian resynchronization in a mouse model of jet lag. *J. Clin. Invest.* 120: 2600–2609.
50. Hut, R. A., A. Scheper, and S. Daan. 2000. Can the circadian system of a diurnal and a nocturnal rodent entrain to ultraviolet light? *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* 186: 707–715.
51. Kenagy, G. J. 1980. Center-of-gravity of circadian activity and its relation to free-running period in two rodent species. *J. Interdiscipl. Cycle Res.* 11: 1–8.
52. Reppert, S. M. 2000. Cellular and molecular basis of circadian timing in mammals. *Semin. Perinatol.* 24: 243–246.
53. Gibson, E. M., C. Wang, S. Tjho, N. Khattar, and L. J. Kriegsfeld. 2010. Experimental “jet lag” inhibits adult neurogenesis and produces long-term cognitive deficits in female hamsters. *PLoS ONE* 5: e15267.
54. Kalsbeek, A., C. X. Yi, C. Cailotto, S. E. la Fleur, E. Fliers, and R. M. Buijs. 2011. Mammalian clock output mechanisms. *Essays Biochem.* 49: 137–151.
55. Boyadjieva, N. I., M. Ortigueira, A. Arjona, X. Cheng, and D. K. Sarkar. 2009. Beta-endorphin neuronal cell transplant reduces corticotropin releasing hormone hyperresponse to lipopolysaccharide and eliminates natural killer cell functional deficiencies in fetal alcohol exposed rats. *Alcohol. Clin. Exp. Res.* 33: 931–937.
56. Dokur, M., N. Boyadjieva, and D. K. Sarkar. 2004. Catecholaminergic control of NK cell cytolytic activity regulatory factors in the spleen. *J. Neuroimmunol.* 151: 148–157.
57. Fu, L., and C. C. Lee. 2003. The circadian clock: pacemaker and tumour suppressor. *Nat. Rev. Cancer* 3: 350–361.
58. Miyazaki, K., M. Wakabayashi, Y. Hara, and N. Ishida. 2010. Tumor growth suppression in vivo by overexpression of the circadian component, PER2. *Genes Cells* 15: 351–358.
59. Ramsey, M. R., and L. W. Ellisen. 2011. Circadian function in cancer: regulating the DNA damage response. *Proc. Natl. Acad. Sci. USA* 108: 10379–10380.
60. Trapani, J. A., and M. J. Smyth. 2002. Functional significance of the perforin/granzyme cell death pathway. *Nat. Rev. Immunol.* 2: 735–747.
61. van den Broek, M. F., D. Kägi, R. M. Zinkernagel, and H. Hengartner. 1995. Perforin dependence of natural killer cell-mediated tumor control in vivo. *Eur. J. Immunol.* 25: 3514–3516.
62. Wallin, R. P., V. Screpanti, J. Michaëlsson, A. Grandien, and H. G. Ljunggren. 2003. Regulation of perforin-independent NK cell-mediated cytotoxicity. *Eur. J. Immunol.* 33: 2727–2735.
63. Davidson, A. J., O. Castanon-Cervantes, T. L. Leise, P. C. Molyneux, and M. E. Harrington. 2009. Visualizing jet lag in the mouse suprachiasmatic nucleus and peripheral circadian timing system. *Eur. J. Neurosci.* 29: 171–180.
64. Yan, L. 2011. Structural and functional changes in the suprachiasmatic nucleus following chronic circadian rhythm perturbation. *Neuroscience* 183: 99–107.
65. Yamazaki, S., R. Numano, M. Abe, A. Hida, R. Takahashi, M. Ueda, G. D. Block, Y. Sakaki, M. Menaker, and H. Tei. 2000. Resetting central and peripheral circadian oscillators in transgenic rats. *Science* 288: 682–685.
66. Kalsbeek, A., I. F. Palm, S. E. La Fleur, F. A. Scheer, S. Perreau-Lenz, M. Ruiters, F. Kreier, C. Cailotto, and R. M. Buijs. 2006. SCN outputs and the hypothalamic balance of life. *J. Biol. Rhythms* 21: 458–469.
67. Hayashi, M., S. Shimba, and M. Tezuka. 2007. Characterization of the molecular clock in mouse peritoneal macrophages. *Biol. Pharm. Bull.* 30: 621–626.
68. Sica, A., L. Dorman, V. Viggiano, M. Cippitelli, P. Ghosh, N. Rice, and H. A. Young. 1997. Interaction of NF-kappaB and NFAT with the interferon-gamma promoter. *J. Biol. Chem.* 272: 30412–30420.
69. Zhou, S., R. Ou, L. Huang, and D. Moskophidis. 2002. Critical role for perforin-, Fas/FasL-, and TNFR1-mediated cytotoxic pathways in down-regulation of antigen-specific T cells during persistent viral infection. *J. Virol.* 76: 829–840.
70. Huang, C., E. Bi, Y. Hu, W. Deng, Z. Tian, C. Dong, Y. Hu, and B. Sun. 2006. A novel NF-kappaB binding site controls human granzyme B gene transcription. *J. Immunol.* 176: 4173–4181.
71. Krukowski, K., J. Eddy, K. L. Kosik, T. Konley, L. W. Janusek, and H. L. Mathews. 2011. Glucocorticoid dysregulation of natural killer cell function through epigenetic modification. *Brain Behav. Immun.* 25: 239–249.
72. Srinivasan, V., G. J. Maestroni, D. P. Cardinali, A. I. Esquifino, S. R. Perumal, and S. C. Miller. 2005. Melatonin, immune function and aging. *Immun. Ageing* 2: 17.
73. Arjona, A., and D. K. Sarkar. 2006. The circadian gene mPer2 regulates the daily rhythm of IFN-gamma. *J. Interferon Cytokine Res.* 26: 645–649.
74. Yang, S., A. Liu, A. Weidenhammer, R. C. Cooksey, D. McClain, M. K. Kim, G. Aguilera, E. D. Abel, and J. H. Chung. 2009. The role of mPer2 clock gene in glucocorticoid and feeding rhythms. *Endocrinology* 150: 2153–2160.
75. Roseboom, P. H., M. A. Nambodiri, D. B. Zimonjic, N. C. Popescu, I. R. Rodriguez, J. A. Gastel, and D. C. Klein. 1998. Natural melatonin “knockdown” in C57BL/6J mice: rare mechanism truncates serotonin N-acetyltransferase. *Brain Res. Mol. Brain Res.* 63: 189–197.
76. Filipiski, E., and F. Lévi. 2009. Circadian disruption in experimental cancer processes. *Integr. Cancer Ther.* 8: 298–302.
77. Innominato, P. F., F. A. Lévi, and G. A. Bjarnason. 2010. Chronotherapy and the molecular clock: Clinical implications in oncology. *Adv. Drug Deliv. Rev.* 62: 979–1001.